

Yes. **Lithium treatment may affect the anion gap**, and in some cases it can cause a **reduced anion gap** or even a **below-zero anion gap**. This happens because lithium is a **positively charged ion** that may not be fully reflected in routine electrolyte panels, so it can shift the balance between measured and **unmeasured electrolytes**. For those interpreting serum chemistry, an **Anion Gap Calculator** can be a practical starting point, but the result always needs the [calculate delta ratio HAGMA](#) broader clinical picture. A medication effect, an electrolyte imbalance, or a broader metabolic disturbance may all be possible explanations.

That said, lithium is only one possible explanation. Proper **laboratory interpretation** requires looking at the full pattern of **electrolytes**, especially **serum sodium**, **blood chloride**, and **bicarbonate**, along with albumin, kidney function, symptoms, and the **lithium concentration**. The anion gap is a useful diagnostic clue, not a standalone diagnosis.

What Is the Anion Gap?

The **anion gap** is a laboratory calculation used to measure the gap between the major measured cations and anions in blood. In practical terms, it helps clinicians evaluate acid-base disorders and spot conditions that may not be obvious from the basic chemistry panel alone. The calculation is based mainly on **sodium**, **chloride**, and **bicarbonate**, all of which are part of routine **serum chemistry**.

In most cases, the body keeps charges balanced. However, not every ion is directly measured. The gap exists because there are **unmeasured ions** in the blood, including proteins, phosphates, sulfates, and certain cations. When the relationship between measured ions changes, the anion gap changes too. That is why it can serve as a clue for **acid-base disorders**, especially **metabolic acidosis**.

One way to understand it: the anion gap compares what is counted in the blood chemistry with what is not counted. If the balance shifts, the gap may become high, low, or, rarely, negative. Changes in **electrolytes** can reflect true disease, medication effects, or a lab issue. That is why interpretation of results should always consider the full clinical setting.


Albumin matters especially because it is one of the main unmeasured anions. When **albumin** is low, the anion gap may fall even if nothing else has changed. That means a low value is not automatically caused by lithium or another toxin.

How an Anion Gap Calculator Works

An **anion gap calculator** is a straightforward utility that runs a **laboratory calculation** using numbers from the electrolyte panel. The most common calculation is:

Anion gap = serum sodium – (serum chloride + serum bicarbonate)

Some tools include potassium, but many do not. The exact formula can vary by lab, so it is important to know which method your clinician or calculator uses. The calculator is only as accurate as the **laboratory values** entered into it, which means mistakes in units, transcription, or specimen quality can affect the result.



When using an anion gap calculator, the main inputs are:

- **Serum sodium**
- **Serum chloride**
- **Serum bicarbonate**

These are the measured ions used to infer the unmeasured portion of the blood's electrical balance. If bicarbonate is low and chloride rises to compensate, the anion gap may appear normal or low depending on the full pattern. If there are extra **positive cations** that are not in the formula, such as lithium, the calculation may show an **anion deficit**.

Because the calculator is based on serum electrolytes, it is not a substitute for clinical judgment. It can quickly flag abnormal lab values, but it cannot determine the cause on its own. The interpretation of results should connect the calculation with symptoms, medication history, and renal function.

Could Lithium Change the Anion Gap?

Indeed. **Lithium** can decrease the anion gap because it acts as an additional **positive cation** in the bloodstream. Since routine chemistry panels do not directly include lithium in the formula, an elevated lithium concentration can reduce the calculated gap. As a result, lithium becomes one of the important **unmeasured cations** that alters the charge balance seen in serum electrolytes.

This is a classic medication effect that can be valuable as a diagnostic clue. If a patient is taking lithium and has a **low anion gap**, clinicians should think about whether the value reflects expected drug exposure, a lab artifact, or **lithium toxicity**. In some cases, the result may even be a **negative anion gap**, although that is rare.

The key mechanism is clear: surplus positively charged lithium ions can decrease the difference between measured sodium and the sum of chloride and bicarbonate. Because the calculation does not count lithium directly, the formula may show a falsely low drop in the gap. This is why lithium is a major consideration in **lab interpretation**, especially when the result does not fit the rest of the clinical picture.

Still, lithium is not the only potential explanation for a low result. The anion gap is influenced by many factors, including albumin, chloride concentration, kidney function, and other causes of metabolic disturbance. That is why the broader context matters more than the number alone.

Reasons Lithium May Lead to a Low Anion Gap

Lithium may produce a low anion gap through its role as an unmeasured cation. In normal acid-base balance, the number of positive and minus charges in the blood is about balanced. If lithium piles up, it introduces extra positive charge without being included in the formula, which can result in **anion gap reduction**. When the effect is marked, the calculated value may become very low or even negative.

This becomes particularly important in **lithium toxicity**. Toxicity can occur when lithium levels rise due to volume depletion, drug interactions, overdose, or impaired kidney clearance. In those cases, the anion gap may offer a clue that something is wrong, but it is not specific enough to confirm the diagnosis by itself. A low gap can point toward medication accumulation, but it can also result from low albumin or analytical issues.

Albumin is important because it is a major unmeasured anion. If albumin is low, the anion gap may already be reduced, and lithium can worsen that effect. That combination can make interpretation more difficult. At the same time, **hyperchloremia** can also push the gap downward, because higher chloride concentration reduces the difference in the formula.

From an acid-base standpoint, a low or negative gap is not the same thing as acidosis, but it can coexist with acid-base problems. That is why clinicians consider both the gap and the broader **acid-base balance**. A patient can have lithium-related changes in the chemistry panel and still also have **metabolic acidosis** from another cause.

In short, lithium may lower the anion gap by adding unmeasured positive charge, but the final result depends on the full metabolic picture.

How to Understand Anion Gap Results in Lithium Users

When an individual on lithium presents with an abnormal anion gap, the result should be handled carefully. The first question is whether the patient has symptoms or signs of **metabolic acidosis**, dehydration, disorientation, shaking, vomiting, or other findings that might suggest toxicity or another acute problem. Then the clinician checks the **lithium level**, because the laboratory result may reflect the drug's actual burden in the body.

Renal function is especially important because lithium is cleared by the kidneys. If kidney function declines, lithium can accumulate, heightening the risk of toxicity and altering the chemistry panel. Even mild **renal impairment** can affect how the body handles lithium, making the medication effect more pronounced.

Clinical interpretation should also include the full electrolyte pattern. For example:

- Reduced sodium with high chloride and low bicarbonate may suggest a different metabolic disturbance.
- A low anion gap with low albumin may reflect hypoalbuminemia rather than lithium.
- A very low or negative value in a patient on lithium should raise concern for toxicity or another lab issue.

A good approach is to compare the anion gap result with the patient's symptoms, medication history, recent dose changes, and kidney function. If the gap is unexpectedly low, repeat testing may help confirm whether the abnormality is real or related to a specimen or calculation problem. This is where careful **lab interpretation** becomes a diagnostic advantage rather than just a numeric exercise.

Also remember that the anion gap is only one part of the assessment. It can be a useful indicator of **electrolyte imbalance**, but lithium users need broader monitoring. When the number does not match the clinical picture, the result should be treated as a clue, not a conclusion.

Additional Sources of an Atypical Anion Gap

It matters not to over-attribute an abnormal anion gap to lithium alone. Numerous conditions can produce a low or high gap, and some can coexist with medication effects. The most common reason for a low gap is **hypoalbuminemia**, because albumin is one of the main unmeasured anions. If albumin is reduced, the apparent gap may fall even when the rest of the chemistry is stable.

Other causes include **hyperchloremia**, analytical variation, and unusual proteins or cations. In the setting of high anion gap metabolic problems, conditions such as **ketoacidosis** and **lactic acidosis** are major considerations. These usually raise the gap rather than lower it, but they are part of the broader differential diagnosis when reading an electrolyte panel.

Toxins can also affect the anion gap. Certain poisonings and medication exposures change measured ions or introduce unmeasured substances that alter the formula. Because the anion gap is a diagnostic clue, not a diagnosis, clinicians use it alongside history, examination, and confirmatory testing.

Other explanations include lab error, changes in protein levels, and shifts in the balance between measured ions and **unmeasured ions**. If the value is very unusual, repeating the blood test is often useful. That helps distinguish a true metabolic disturbance from an artifact.

Ultimately, a low or abnormal gap should be interpreted in context. Lithium is on the list, but so are albumin changes, kidney problems, acid buildup, and toxic exposures.

When to Request Medical Evaluation

Clinical review is important whenever an abnormal anion gap appears together with concerning **symptoms**. This is especially true for someone taking lithium. Signs such as vomiting, diarrhea, confusion, severe tremor, unsteady walking, marked drowsiness, or worsening weakness can signal the need for urgent assessment. These symptoms may indicate **lithium toxicity** or another serious metabolic issue.

Immediate review is recommended when signs are serious, the patient looks different, or there is worry for overdose, dehydration, or kidney failure. Because lithium can become risky when levels go up, prompt assessment is essential. A clinician may order follow-up electrolytes, a **lithium level**, kidney function tests, and other studies depending on the clinical picture.

Drug level monitoring is a key part of lithium therapy. In case the anion gap is abnormal, the medication dose, recent changes, hydration status, and other medications should be reviewed. Numerous interacting drugs can affect kidney clearance and increase risk. The clinician may also look for **renal function** changes that could explain the abnormal lab value.

A doctor's evaluation is the appropriate next step whenever the lab result is unexpected or the patient does not feel well. The goal is to separate a benign variation from a genuine clinical concern. In reality, that means pairing the chemistry panel with symptoms, medication effect, and diagnostic testing rather than relying on one number alone.

FAQ: Lithium and Anion Gap Questions

Can lithium cause a low anion gap?

Yes. Lithium can cause a **low anion gap** because it is a **charged cation** that is not included in the typical anion gap formula. When lithium is present in sufficient amounts, it can reduce the calculated gap by adding

unmeasured cations to the blood chemistry profile.

Can lithium lead to a below-zero anion gap?

Certainly, while it is rare. A **negative anion gap** can happen when lithium accumulation is substantial or when other factors, like low albumin or high chloride, also shift the calculation down. A negative result should warrant careful examination of the labs and clinical status.

In what way does an anion gap calculator help interpret lithium-related lab results?

An **anion gap calculator** offers a quick **laboratory calculation** from **serum sodium**, **serum chloride**, and **serum bicarbonate**. For people taking lithium, it can point to an unexpected **anion deficit** that may be a sign to medication effect, toxicity, or another metabolic disturbance. It can assist, but does not replace, clinical interpretation.

Does a low anion gap necessarily mean lithium?

No. **Hypoalbuminemia** is a very common cause of a low gap, and **hyperchloremia**, lab variation, and other conditions can also lower it. Lithium is an important possibility, but it should not be assumed to be the only explanation without looking at albumin, kidney function, and the rest of the electrolytes.

What needs to be checked if someone on lithium has an abnormal anion gap?

Clinicians usually check the **lithium level**, **renal function**, albumin, repeat electrolytes, and the full clinical picture. They also assess for **symptoms** of toxicity or other acid-base disorders such as **metabolic acidosis**. This combined approach improves **lab interpretation** and helps determine whether the abnormal value reflects a true problem or a transient finding.