



HOW WE TREAT CAR ACCIDENT INJURIES IN LAKEWOOD

Every recovery plan at Injury Recovery Center is built around your body, your injury, and your goals—not a template.



MANUAL CHIROPRACTIC ADJUSTMENTS

Targeted, hands-on treatment focused on restoring movement and reducing pain. We don't use "clicker" tools or drop tables. No aggressive "grip it and rip it" adjustments. Just be careful, intentional care that progresses at your pace.



INTERSEGMENTAL TRACTION THERAPY

Gentle traction therapy that supports spinal mobility and helps break down scar tissue before it hardens into chronic restriction.



SHOCKWAVE THERAPY

Non-invasive treatment that activates your body's natural healing response for persistent soft tissue pain and inflammation. Non-treatment that activates your body's natural healing response for persistent soft tissue pain and inflammation.



THERAPEUTIC EXERCISES & MODALITIES

Customized rehabilitation exercises are designed to restore strength and stability after trauma.



MEDICAL & SPECIALIST REFERRALS

When your recovery requires care beyond chiropractic—physical therapy, orthopedics, neurology, pain management, or mental health support—we coordinate those referrals. Dr. Ron operates as a hub in a medical-legal network, not a silo.



Whiplash is often framed as a neck problem, which is understandable. The pain is usually felt in the neck first. The stiffness lives there. Headaches often start there. Turning the head while driving or looking down at a phone can suddenly feel complicated. But anyone who has worked with people recovering from whiplash for any length of time learns the same lesson: the neck rarely suffers alone.

The body responds to a sudden acceleration-deceleration injury as a connected system. The jaw may tighten. The shoulders brace. Breathing becomes shallow. The ribs lose motion. The low back may ache, even if it did not hit anything. Balance changes. Sleep gets worse. Then a pattern develops. A person who was injured in a car accident starts moving less, guarding more, and using the trunk in a rigid, inefficient way. That is where core strength enters the picture, not as a fitness trend, but as part of sound whiplash therapy.

A strong, responsive core does not “fix” whiplash by itself. That would be too simple, and frankly, misleading. What it can do is create the conditions for better movement, reduced strain, improved tolerance for daily tasks, and a more durable recovery. The key is knowing what “core” really means and how to train it without provoking symptoms.

The neck does not work in isolation

When people hear the phrase core strength, they often picture six-pack abs or intense gym work. In rehabilitation, the term means something much broader. It includes the deep and superficial muscles that help control the spine, ribs, pelvis, and pressure within the trunk. It also includes timing. A person can be very strong in a gym sense and still have poor trunk control after injury.

That matters because the neck sits on top of a moving base. If the rib cage is stiff, the shoulder blades are poorly controlled, and the upper back does not rotate well, the neck has to do extra work. If the trunk is unstable or excessively rigid, the small muscles around the cervical spine often step in and overwork. Patients describe this in plain language. “My neck gets tired just sitting up.” “I can’t hold my posture by the afternoon.” “If I carry groceries, my headache starts.” Those are not always signs of a neck that is weak in isolation. They often reflect a body that has lost coordination.

After whiplash, it is common to see an altered pattern of movement. The person may move as one block instead of segmentally. They may stop rotating through the thoracic spine and compensate with the neck. Their breathing may become chest-dominant, with overactivity in the upper traps and accessory breathing muscles. The pelvis may become fixed. Walking may shorten. The nervous system becomes protective, and that protective strategy, useful in the first days after injury, can become a problem if it lingers for weeks or months.

This is why effective whiplash therapy usually expands beyond the painful area. A neck-only program can miss the mechanical drivers that keep symptoms going.

What actually happens after whiplash

Whiplash is not one uniform injury. Severity varies. So does tissue involvement. Some people recover quickly within a few weeks. Others develop a much longer course with persistent pain, dizziness, headaches, jaw symptoms, concentration difficulty, exercise intolerance, and fear of movement.

The mechanism matters. A rear-end collision at low to moderate speed can still produce significant symptoms, especially if the person was not prepared for impact. The body is taken through a rapid sequence it did not organize for. Soft tissues can be strained. denvercarcrashdoctor.com Whiplash Therapy Joints can become irritated. Muscles can guard. The nervous system can become sensitized. Stress and poor sleep can amplify the whole picture.

What tends to surprise people is how often the symptoms spread beyond the neck. Mid-back pain is common. Low back discomfort is not unusual. Some people feel unstable when walking in crowded places or turning quickly. Others notice that their symptoms spike after sitting for long periods rather than after obvious exertion. This is one reason simplistic advice, such as “just rest your neck,” often falls short after the acute stage.

Rest has a place early on, especially in the first days when pain is sharp and movement feels threatening. But prolonged withdrawal from movement usually backfires. Tissues decondition. The trunk gets less supportive. The breathing pattern becomes less efficient. The person grows less confident. Then ordinary life feels like a workout.

Why core strength matters more than most people realize

The most practical way to think about core strength in whiplash recovery is as load sharing. When the trunk can absorb and transfer force well, the neck does not have to manage every demand on its own.

Imagine a person reaching into the back seat of a car, lifting a toddler, or carrying a laptop bag up stairs. Those tasks are not neck exercises, but they challenge the entire system. If the lower ribs flare, the pelvis tips forward, and the upper traps tighten to create stability, the neck often pays the price. If the trunk can stay quietly organized, the movement becomes less threatening and more efficient.

There is also a timing issue. Healthy movement is full of subtle anticipatory adjustments. Before the arm lifts, parts of the trunk prepare. Before a quick turn, the body sequences rotation. After whiplash, this timing can become delayed or overly rigid. People “hold” themselves upright instead of being supported dynamically. They are often exhausted by posture because posture has turned into effort.

A stronger, better-coordinated core can help with:

- reducing unnecessary strain on the cervical muscles during daily tasks
- improving postural endurance without forcing a stiff upright position
- supporting better breathing mechanics, which can lower tension and improve exercise tolerance
- restoring smoother rotation through the thoracic spine and trunk
- increasing confidence with lifting, carrying, walking, and returning to work

Notice the word coordinated. Many patients have already tried generic abdominal exercises by the time they seek more formal help. Sit-ups, [Whiplash Therapy](#) long planks, or intense bootcamp classes can actually aggravate symptoms if introduced too early or without context. The goal is not to brace harder. It is to create support without excessive guarding.

Breathing, ribs, and the hidden burden on the neck

One of the most overlooked pieces of whiplash therapy is breathing. Not because breathing is mystical, but because the mechanics matter. After neck injury, many people begin to breathe high in the chest. Their scalenes, upper trapezius, and sternocleidomastoid become overused. Every breath then reinforces neck tension.

If the diaphragm is not doing its share, the trunk loses one of its most important stabilizing contributors. The ribs stay elevated. The neck muscles become both movers and stabilizers. That is an expensive strategy. It often drives headaches, neck fatigue, and a sense that one can never fully relax.

A good therapist will often watch a patient breathe before prescribing much exercise. Are the lower ribs moving? Can the patient exhale without collapsing? Does inhalation create neck tension? Can they maintain an easy breath while moving an arm or leg? These are not trivial details. They are often the bridge between symptom relief and symptom recurrence.

I have seen patients whose pain flared every time they tried “core work,” only to improve once the exercises were reframed around breath, rib positioning, and low-level control. In practical terms, that may mean starting with

supported positions, very short holds, or movements that feel almost too easy at first. The point is not to impress anyone. The point is to rebuild function that transfers into daily life.

The difference between fitness core work and rehab core work

This distinction is worth making clearly. Traditional fitness training often emphasizes challenge, fatigue, and progressive overload. Rehabilitation after whiplash emphasizes symptom response, precision, and tolerance.

In a gym setting, a coach might chase longer planks, heavier carries, or high-rep abdominal work. In a whiplash setting, that can be too much, too soon. A patient may compensate by clenching the jaw, elevating the shoulders, or holding the breath. The exercise looks successful from the outside but leaves them with a headache that night.

Rehabilitation usually starts with quieter tasks. Can the patient lie comfortably and breathe without neck gripping? Can they gently engage the trunk while moving a limb? Can they roll, sit, stand, and turn with less stiffness? Can they walk briskly for ten minutes without a symptom spike later in the day? Those wins may seem modest, but they are often exactly what lets someone return to driving, desk work, parenting, or light training.

Later, if recovery is progressing well, the work can become more demanding. Carries, resisted rotation, controlled hinging, step patterns, and whole-body strengthening all have value. The order matters. The dose matters. The body's response matters even more.

Why some people get stuck

Not every slow recovery comes down to poor core strength. That would be an oversimplification. Persistent symptoms can involve joint irritation, vestibular issues, visual sensitivity, concussion-related concerns, jaw dysfunction, stress physiology, poor sleep, central sensitization, and workplace demands. Whiplash therapy often works best when it respects that complexity rather than pretending there is one silver bullet.

Still, there are predictable reasons people stall.

Sometimes the early advice was too passive. They were told to rest, use a collar, and wait. Weeks passed, movement shrank, and the body became less adaptable.

Sometimes they were pushed too hard. They were handed generic strengthening drills or told to "push through it," which reinforced flares and fear.

Sometimes the neck was treated in isolation. Manual therapy may have helped temporarily, but no one addressed trunk control, breathing, or movement habits.

And sometimes the patient did everything right but had a nervous system that remained highly protective. In those cases, the work often requires more patience, smaller doses, and better pacing.

That last point deserves emphasis. Progress in whiplash recovery is not always linear. A person may tolerate walking but struggle with sitting. They may handle exercise but flare after a stressful workday. They may feel much better, then have a setback after travel or a poor night of sleep. Those fluctuations do not necessarily mean the program is failing. They usually mean the system still has a limited buffer.

What good whiplash therapy tends to include

There is no universal formula, but effective care often blends several elements rather than relying on one technique. The best programs are adaptable. They respect irritability, but they do not let fear dictate the entire plan.

A thoughtful clinician may include the following:

- education about pain, pacing, and realistic recovery patterns
- gentle neck mobility and deep neck flexor retraining when appropriate
- breathing and rib mechanics work to reduce neck overuse
- thoracic mobility and shoulder girdle control
- graded trunk and whole-body strengthening that reflects real life demands

This is where clinical judgment matters. A patient with dizziness may need a different sequence than a patient whose main issue is neck fatigue at a desk. A new parent lifting an infant repeatedly needs different task practice than a construction worker returning to overhead labor. The principles overlap, but the expression should fit the person.

The office worker and the athlete are not the same case

Two people can carry the same diagnosis and need very different rehabilitation.

Take the office worker with headaches that build after two hours at a screen. Their rehab may hinge on breathing, visual breaks, upper thoracic mobility, desk positioning, and improved trunk support in sitting and standing transitions. If you only give them neck stretches, you may miss the bigger driver, which is low-grade, sustained strain through the entire system.

Now consider the recreational tennis player who feels fine at rest but develops neck pain when serving or rotating quickly. That case may demand more dynamic trunk rotation, deceleration control, scapular timing, and confidence under speed. Static postural drills alone will not prepare them for sport.

Or think about the patient with both neck pain and low back pain after a crash. This is common enough to deserve attention. If the lumbar spine is sore and the person avoids bending, they often become even more rigid through the trunk. Then every turn gets shunted upward. In that situation, core strength is not optional background work. It is part of restoring the ability to move without bracing from the jaw down.

How to load the body without feeding symptoms

One of the hardest parts of whiplash therapy is dosing. Too little challenge and recovery drifts. Too much and the patient flares. A useful clinical rule is that symptoms during exercise are not automatically bad, but they should be understandable, limited, and settle within a reasonable window. For many patients, a mild increase that eases within hours is acceptable. A flare that ruins sleep or worsens the next day's baseline usually means the dose was too high.

This is where people benefit from specific coaching. They need to learn the difference between effort and threat. Muscles working is fine. Holding the breath, clenching the jaw, and finishing every session with a pounding headache is not.

A few practical markers help guide progression:

- symptoms stay stable or only mildly increase during activity
- breathing remains smooth rather than shallow and urgent
- the neck does not take over every movement
- recovery after the session is predictable
- confidence in daily tasks gradually improves

Those markers are often more useful than chasing arbitrary repetitions.

The psychology of support

There is another layer to core work that does not get discussed enough. When people regain control through the trunk, they often feel safer in their own body. That matters. Whiplash can make ordinary motion feel unreliable. People stop trusting turns, sudden stops, lifting, or exercise. They begin to monitor every sensation. A body that feels supported, responsive, and less fragile changes that relationship.

This is not simply mental. It is embodied. When someone learns they can carry bags, rotate to reverse the car, or walk uphill without paying for it later, fear tends to drop. That reduction in threat can lower guarding, which further improves movement. It becomes a positive cycle.

The reverse is also true. If rehabilitation is framed in a way that makes the person feel damaged, unstable, or permanently vulnerable, the body often tightens around that story. Good therapy is not false reassurance, but it should leave room for capability.

Where manual therapy fits, and where it does not

Hands-on treatment can be helpful. Soft tissue work, gentle mobilization, and symptom-modifying techniques can reduce pain enough for someone to move better. But manual therapy is usually a door, not the entire house.

If a patient feels looser for six hours and then reverts to the same movement pattern, the underlying problem remains. The body still needs a reason to organize differently. That reason often comes from exercise, graded exposure, breathing work, and repeated practice in meaningful tasks.

This is especially true for core-related deficits. No amount of massage can build endurance for sitting upright through a work meeting or carrying groceries up two flights of stairs. Relief matters, but function is the long game.

When to be cautious

Most people with whiplash improve with conservative care, but caution is appropriate when symptoms are severe, progressive, or unusual. Significant neurological signs, marked weakness, worsening numbness, unexplained fainting, severe unremitting headache, or signs of more serious injury deserve medical evaluation. Persistent dizziness, vision changes, or concussion-related symptoms may require a broader team approach.

It is also wise to be careful with aggressive exercise in the early phase if even small loads trigger major flares. Those cases often respond better to reduced complexity, lower intensity, and tighter pacing before progressing.

None of this argues against strengthening. It argues for matching the plan to the presentation.

The body matters because recovery is lived, not isolated

People do not recover from whiplash in a vacuum. They recover while commuting, parenting, working, exercising, sleeping badly, and trying to feel normal again. The neck is part of that reality, but not the whole of it.

Core strength, used in the rehabilitation sense, helps reconnect the body into something useful and dependable. It gives the neck a better foundation. It restores load sharing. It improves breathing mechanics. It supports posture without turning posture into strain. It builds tolerance for the ordinary tasks that tell a person they are getting their life back.

That is why good Whiplash Therapy pays attention to the entire body. Not because every patient needs a complicated program, and not because core work is fashionable, but because the body is how recovery happens. When treatment respects that, progress tends to be more durable, more practical, and more honest.

Injury Recovery Center

2290 Kipling St Unit 6

Lakewood, CO 80215, United States

Phone: +1 720-575-8791

FAQ About Whiplash Therapy

How long does it take for a neck whiplash to heal?

Many people improve within a few weeks to two or three months, but symptoms can last longer. Recovery depends on the injury and individual circumstances. A clinician can assess persistent symptoms and adjust care as needed; a fixed recovery date cannot be guaranteed.

What not to do after whiplash?

Avoid prolonged bed rest, forceful neck movements, and wearing a neck collar unless a clinician recommends it. Follow advice about gentle activity and exercise. New weakness, tingling, difficulty walking, or severe worsening pain needs prompt medical evaluation.

How to heal whiplash at home?

After an appropriate assessment, home care may include gentle everyday activity, prescribed movement exercises, and heat or cold for comfort. Ask a clinician or pharmacist which pain medicines are safe for you. Seek reassessment if symptoms persist or worsen; home care does not replace evaluation of a potentially serious neck injury.